# Equi-rank Homomorphism Preservation Theorem on Finite Structures

Benjamin Rossman Duke University

FMT 2025, Les Houches

preserved under *injective / surjective / all* homomorphisms if, and only if, it is

equivalent to an *existential / positive / existential-positive* sentence.

preserved under *injective / surjective / all* homomorphisms if, and only if, it is

equivalent to an *existential / positive / existential-positive* sentence.

- Originally proved in the 1950's using **compactness**.
- Proofs give **no computable upper bound** on the quantifier rank of the equivalent existential / positive / existential-positive sentence.
- Proofs relativize to any FO-axiomatizable class of structures.
   Proofs do not relativize to the non-FO-axiomatizable class of *finite* structures.

preserved under *injective / surjective / all* homomorphisms

if, and only if, it is

equivalent to an *existential / positive / existential-positive* sentence.

**Question** [Gurevich '84] What is the status of classical theorems when restricted to *finite structures*?

preserved under *injective / surjective / all* homomorphisms if, and only if, it is

equivalent to an *existential / positive / existential-positive* sentence.

Question [Gurevich '84]

What is the status of classical theorems when restricted to *finite structures*?

- $\bullet$  Compactness <u>fails</u> on finite structures.
- The Los-Tarski and Lyndon Preservation Theorems <u>fail</u> on finite structures [Tait '59, Ajtai-Gurevich '87].
- In contrast, the **HPT** is <u>valid</u> on finite structures [R.'05].

Every is equivalent to an hom-preserved sentence with quant. rank r existential-positive sentence with quant. rank r

#### HPT on Finite Structures [R.'05]

Every is equivalent to an hom-preserved sentence with quant. rank r existential-positive sentence with quant. rank r

#### HPT on Finite Structures [R.'05]

## Equi-rank HPT on Finite Structures [R.'25]

<u>On finite structures</u>, every is equivalent to an hom-preserved sentence with quant. rank r existential-positive sentence with quant. rank r

Every is equivalent to an hom-preserved sentence with quant. rank r existential-positive sentence with quant. rank r

• MAIN LEMMA

Every structure  $\mathfrak{A}$  has an (infinite)  $\exists^+$ -saturated co-retract  $\widehat{\mathfrak{A}} \xrightarrow{\supset} \mathfrak{A}$  with the property that

$$\mathfrak{A} \equiv^{\exists^+}_r \mathfrak{B} \implies \widehat{\mathfrak{A}} \equiv^{\mathsf{FO}}_r \widehat{\mathfrak{B}}$$

Every is equivalent to an hom-preserved sentence with quant. rank r existential-positive sentence with quant. rank r

## HPT on Finite Structures [R.'05]

• MAIN LEMMA

Every <u>finite</u> structure  $\mathfrak{A}$  has an <u>finite</u> co-retract  $\widehat{\mathfrak{A}}_r$ , with the property that

$$\mathfrak{A} \equiv_{f(r)}^{\exists^+} \mathfrak{B} \implies \widehat{\mathfrak{A}}_r \equiv_r^{\mathsf{FO}} \widehat{\mathfrak{B}}_r$$

• REMARK

The  $\exists^+$ -saturated co-retract  $\widehat{\mathfrak{A}}$  (in the equi-rank proof) is a direct limit

 $\mathfrak{A} = \widehat{\mathfrak{A}}_0 \xleftarrow{\subset} \widehat{\mathfrak{A}}_1 \xleftarrow{\subset} \widehat{\mathfrak{A}}_2 \xleftarrow{\subset} \cdots \xleftarrow{\subset} \widehat{\mathfrak{A}}$ 

Every is equivalent to an hom-preserved sentence with quant. rank r existential-positive sentence with quant. rank r

## HPT on Finite Structures [R.'05]

Improvements to  $f : \mathbb{N} \to \mathbb{N}$ 

$$2005 \quad f(r) = 2^{2^{2^{-1}}} \quad \text{co-retracts } \widehat{\mathfrak{A}}_r \xrightarrow{\supset} \mathfrak{A}$$

$$2017 \quad f(r) = 2^{O(r)}$$

$$2018 \quad f(r) = \widetilde{O}(r^5)$$

$$2019 \quad f(r) = \widetilde{O}(r^3)$$

$$2025 \quad f(r) = r \quad [R.'05] + \text{Cai-Fürer-Immerman construction}$$

## *Exponential-rank* HPT on Finite Structures [R.'05] + [R.'14, R.'17]

- Combines the previous result with a tight lower bound on the  $AC^0$  formula size (= FO[Arb] quantifier rank) of DISTANCE-k CONNECTIVITY.
- Uses the fact that the **longest path** in a graph gives an *exponential approximation* of tree-depth:  $\log(lp(G)) \leq td(G) \leq lp(G)$

## *Exponential-rank* HPT on Finite Structures [R.'05] + [R.'14, R.'17]

**Poly-rank HPT on Finite Structures** [R.'05] + [Li-Razborov-R.'14, R'17, Kawarabayashi-R.'18, Kush-R.'20, Czerwinski-Nadara-Pilipczuk '19]

• Relies on a *polynomial excluded-minor approximation* of **treedepth** (in terms of **longest path**, tallest complete **binary tree minor**, and **tree-width**).

## *Exponential-rank* HPT on Finite Structures [R.'05] + [R.'14, R.'17]

**Poly-rank HPT on Finite Structures** [R.'05] + [Li-Razborov-R.'14, R'17, Kawarabayashi-R.'18, Kush-R.'20, Czerwinski-Nadara-Pilipczuk '19]

<u>On finite structures</u>, every hom-preserved FO[Arb] sentence with quant. rank r is equivalent to an existential-positive sentence with quant. rank  $\widetilde{O}(r^3)$ 

#### Equi-rank HPT on Finite Structures [R.'25]

• Proof combines the original **HPT on Finite Structures** [R.'05] with a generalized **Cai-Fürer-Immerman construction** [CFI '94].

#### **Generalized Cai-Fürer-Immerman Construction**

Let  $\mathfrak{A}$  be a finite structure with universe V, and let  $\alpha : V \to \mathbb{Z}_2$ .

Then  $\mathbf{CFI}_{\alpha}(\mathfrak{A})$  is a structure—in the same relational signature—whose elements are pairs  $(v, \lambda)$  such that  $v \in V$  and  $\lambda : \mathrm{Nbd}(v) \to \mathbb{Z}_2$  and

$$\sum_{w \in \mathrm{Nbd}(v)} \lambda(w) = \alpha(v).$$

For each *t*-ary relation  $R \subseteq V^t$  in  $\mathfrak{A}$ , the corresponding relation in  $\mathbf{CFI}_{\alpha}(\mathfrak{A})$  consists of *t*-tuples  $((v_1, \lambda_1), \ldots, (v_t, \lambda_t))$  such that  $(v_1, \ldots, v_t) \in R$  and

$$\lambda_i(v_j) = \lambda_j(v_i)$$
 for all  $i, j \in \{1, \dots, t\}$ .

OBS: The projection  $(v, \lambda) \mapsto v$  is a homomorphism  $\mathbf{CFI}_{\alpha}(\mathfrak{A}) \to \mathfrak{A}$ .

#### **Generalized Cai-Fürer-Immerman Construction**

Let  $\mathfrak{A}$  be a finite structure with universe V, and let  $\alpha : V \to \mathbb{Z}_2$ .

Then  $\mathbf{CFI}_{\alpha}(\mathfrak{A})$  is a structure—in the same relational signature—whose elements are pairs  $(v, \lambda)$  such that  $v \in V$  and  $\lambda : \mathrm{Nbd}(v) \to \mathbb{Z}_2$  and

$$\sum_{w \in \mathrm{Nbd}(v)} \lambda(w) = \alpha(v).$$

For each *t*-ary relation  $R \subseteq V^t$  in  $\mathfrak{A}$ , the corresponding relation in  $\mathbf{CFI}_{\alpha}(\mathfrak{A})$  consists of *t*-tuples  $((v_1, \lambda_1), \ldots, (v_t, \lambda_t))$  such that  $(v_1, \ldots, v_t) \in R$  and

$$\lambda_i(v_j) = \lambda_j(v_i)$$
 for all  $i, j \in \{1, \dots, t\}$ .

OBS: The projection  $(v, \lambda) \mapsto v$  is a homomorphism  $\mathbf{CFI}_{\alpha}(\mathfrak{A}) \to \mathfrak{A}$ .

#### Key Lemma

Suppose  $\mathfrak{A}$  is a core (i.e. every homomorphism  $\mathfrak{A} \to \mathfrak{A}$  is an isomorphism).

Then there exists a homomorphism  $\mathfrak{A} \to \mathbf{CFI}_{\alpha}(\mathfrak{A})$  if, and only if,  $\sum_{u \in U} \alpha(u) = 0$  for every connected component  $U \subseteq V$  (in the Gaifman graph of  $\mathfrak{A}$ ).

#### **Generalized Cai-Fürer-Immerman Construction**

Well-known property of CFI structures [CFI '94]

If  $\mathfrak{A}$  has **tree-depth** > r (respectively, **tree-width** > k), then for all  $\alpha, \beta$  :  $V \to \mathbb{Z}_2$ , structures  $\mathbf{CFI}_{\alpha}(\mathfrak{A})$  and  $\mathbf{CFI}_{\beta}(\mathfrak{A})$  are indistinguishable by first-order sentences with **quantifier rank** r (respectively, **variable width** k).

#### Equi-rank HPT on Finite Structures [R.'25]

Suppose  $\Phi$  is first-order sentence of quantifier rank r that is preserved under homomorphisms on finite structures. Then  $\Phi$  is equivalent on finite structures to an  $\exists^+$  sentence of quantifier rank r.

## PROOF

• By the **HPT on Finite Structures**, the class  $Mod_{fin}(\Phi)$  is generated by finitely many  $\rightarrow$ -incomparable *minimal cores*  $\mathfrak{C}_1, \ldots, \mathfrak{C}_m$ . That is, for all finite structures  $\mathfrak{A}$ ,

$$\mathfrak{A} \models \Phi \iff \bigvee_{i=1}^{m} (\mathfrak{C}_i \to \mathfrak{A}).$$

• By lemmas in [R.'05], it suffices to show that each  $\mathfrak{C} \in {\mathfrak{C}_1, \ldots, \mathfrak{C}_m}$  has **tree-depth**  $\leq r$ .

#### Equi-rank HPT on Finite Structures [R.'25]

Suppose  $\Phi$  is first-order sentence of quantifier rank r that is preserved under homomorphisms on finite structures. Then  $\Phi$  is equivalent on finite structures to an  $\exists^+$  sentence of quantifier rank r.

# **PROOF** (continued)

- <u>CLAIM</u>: Every *minimal core*  $\mathfrak{C} \in Mod_{fin}(\Phi)$  has tree-depth  $\leq r$ .
- Let V be the universe of  $\mathfrak{C}$  and let even, odd :  $V \to \mathbb{Z}_2$  such that

even  $\equiv 0$  and  $\sum_{v \in V} \operatorname{odd}(v) = 1$ .

- By the **Key Lemma**,  $\mathfrak{C} \to \mathbf{CFI}_{\mathsf{even}}(\mathfrak{C})$  and  $\mathfrak{C} \not\to \mathbf{CFI}_{\mathsf{odd}}(\mathfrak{C}) \to \mathfrak{C}$ .
- Therefore,  $\mathbf{CFI}_{\mathsf{even}}(\mathfrak{C}) \models \Phi$  (since  $\Phi$  is hom-preserved) and  $\mathbf{CFI}_{\mathsf{odd}}(\mathfrak{C}) \not\models \Phi$  (since  $\mathfrak{C}$  is a minimal core).
- Since  $\mathbf{CFI}_{\mathsf{even}}(\mathfrak{C})$  and  $\mathbf{CFI}_{\mathsf{odd}}(\mathfrak{C})$  are distinguished by a first-order sentence of quantifier rank r, it follows that  $\mathfrak{C}$  has  $\mathbf{tree-depth} \leq r$  (by the well-known property of CFI structures).

# Q.E.D.

A refinement of this argument establishes a stronger version of our equi-rank result in terms of multiple syntactic parameters:

#### **Equi-Resource HPT on Finite Structures** [R.25]

<u>On finite structures</u>, every homomorphism-preserved sentence is equivalent to an existential-positive sentence with the same *quantifier rank*, *number of variables*, and *alternation type*.